



Oesophago-Gastric Anastomosis Audit Protocol 2018

www.ogaa.org.uk

Email: OGanastomosisaudit@gmail.com

 [@OGAAudit](https://twitter.com/OGAAudit)

Steering Group

Name	Organisation	Email
Mr Richard Evans	General Surgery Trainee – West Midlands Deanery	rptevans@doctors.org.uk
Mr Pritam Singh	General Surgery Trainee – West Midlands Deanery	pritam@cantab.net
Mr Sivesh Kamarajah	University of Birmingham	SXK206@student.bham.ac.uk
Mr James Bundred	University of Birmingham	JXB473@student.bham.ac.uk
Dr Dmitri Nepogodiev	University of Birmingham	D.Nepogodiev@bham.ac.uk
Mr Benjamin Jefferies	University of Birmingham	bjj416@student.bham.ac.uk
Mr Kwabena Siaw-Acheampong	University of Birmingham	kxs598@student.bham.ac.uk
Ms Siobhan McKay	General Surgery Trainee – West Midlands Deanery	mckay.siobhan@gmail.com
Mr Kasun Wanigsooriya	General Surgery Trainee – West Midlands Deanery	kasun87@live.co.uk
Dr Tony Whitehouse	University Hospitals Birmingham NHS Foundation Trust	Tony.Whitehouse@uhb.nhs.uk
Prof Derek Alderson	Royal College of Surgeons England	dalderson@rcseng.ac.uk
Mr James Gossage	Guy's and St. Thomas' NHS Foundation Trust	james.gossage@gstt.nhs.uk
Prof Richard van Hillegersberg	UMC Utrecht	R.vanHillegersberg@umcutrecht.nl
Mr Ravinder Vohra	Nottingham University Hospitals NHS Trust	Ravinder.Vohra@nuh.nhs.uk
Mr Ewen Griffiths	University Hospitals Birmingham NHS Foundation Trust	ewen.griffiths@uhb.nhs.uk

Supporting organisations

West Midlands Surgical Research Collaborative



Birmingham Clinical Trials Unit

Providing REDCap Access



UNIVERSITY OF
BIRMINGHAM

Academic Department of Surgery



Association of Laparoscopic Surgeons GB&I

AUGIS

Association of Upper Gastrointestinal Surgeons of
Great Britain and Ireland

Lay Summary

Oesophageal cancer is the sixth leading cause of cancer related death affecting up to 450,000 people globally each year. The main surgical treatment for oesophageal cancer is oesophagectomy - an operation to remove part of the oesophagus and stomach followed by a join between the remaining oesophagus and stomach. The techniques used to create this join vary and involve various stitching methods and stapling devices. A proportion of these joins will breakdown and this can result in the patients becoming very unwell with a risk of death. The strategies to manage this complication also vary and include:

- No surgical intervention
- An endoscopic intervention or
- A further surgical procedure.

This international audit will look at the rates of breakdown of these joins, commonly termed a 'leak', how they are managed and the effect on the patient outcomes. The information collected from this audit will help to develop recommendations on how to prevent and manage this serious complication.

Short Summary

Primary Audit Objectives

- 1- Quantify the incidence of anastomotic leak rate in an international multicentre audit which incorporates data from high and low volume centres and high and low income countries
- 2- Assess the variation in anastomotic leak rates internationally
- 3- Assess the relationship between anastomotic technique and optimal patient outcome – discharge home eating and drinking orally
- 4- Assess the relationship between anastomotic leak therapy and optimal patient outcome

Audit Standard

- 1- Anastomotic leak and conduit necrosis rate should be less than 13%
- 2- Major post-operative morbidity (Clavien Dindo Grade III or more) should be less than 35%.
- 3- 30 day mortality rate should be less than 5% and 90 day mortality rate should be less than 8%.

A data collection protocol will identify patient demographics, operative and peri-operative details and outcome markers. Key outcome measures will include post-operative mortality, morbidity including grade of leak and length of stay. Management techniques used for anastomotic leaks will also be assessed (e.g. conservative management, oesophageal stent, endo-luminal VAC therapy and re-operation).

Methods

A nine month multicentre prospective audit will be performed globally starting in April 2018 and coordinated by University Hospitals Birmingham. This will include patients undergoing oesophagectomy over 6 months and encompassing a 90-day follow up period. A pilot data collection period will occur at University Hospitals Birmingham and 3 other UK hospitals in 2017. Sites will be required to pre-register for the audit and obtain local study approval prior to commencement of the study.

During the study sites will be required to record data contemporaneously via a dedicated encrypted server through the Research Electronic Data Capture (REDCap) web application secure online database. The REDCap database will provide a standardised data collection proforma assessing key information to answer the primary audit question. The report of the audit will be prepared in accordance with the guidelines as set by the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) statement for observational studies and STROCSS (Strengthening the Reporting of Cohort Studies in Surgery)(1). All unit results will be anonymised to all but the auditors and the specific unit. Unit results will not be shared with other units or the collaborators as a whole. The study will be defined as audit not research in accordance with the NHS Health research authority recommendations (Appendix 2).

Discussion

Data for this multicentre international audit will be collected by both surgeons and trainees to provide greater insight into the complexities of oesophagectomy and outcome. This audit may highlight trends in improved survival associated with specific operative techniques or specific management strategies to deal with leaks that can be further assessed and analysed through research to improve outcomes in oesophageal cancer.

Introduction

Oesophageal cancer is the sixth leading cause of cancer related mortality affecting up to 450,000 people globally each year. There are 2 histological types – squamous cell carcinoma and adenocarcinoma. While the incidence of squamous cell cancer is stable worldwide, the incidence of adenocarcinoma has risen dramatically in the western world in parallel with obesity over the last 30 years. Despite advances in modern treatment, 5-year survival remains at around 15 to 20%. Oesophagectomy is a mainstay in curative treatment for those with oesophageal cancer however the technique and outcome varies greatly. The techniques used for oesophagectomy vary greatly, particularly regarding operative technique and methods of reconstruction.

Anastomotic leakage following oesophagectomy is one of the most feared complications associated with high rates of morbidity and mortality. 30 day mortality in patients with a demonstrable leak can be as high as 17-35% whereas the 30 day mortality of patients with an intact anastomosis is 2-3% (2, 3). In addition anastomotic leakage is known to increase length of hospital stay, reduced quality of life and be economically costly for the health service(4). There is also evidence that anastomotic leakage affects long term prognosis and is associated with reduced long term survival and increased recurrence rates (5).

Anastomotic leak rates are very variable between surgeons, units and countries. Current practices demonstrate rates between 1.8% and 18.2% (6-15). The largest of the recent studies by Kassis et al identified 7,595 oesophagectomies with a leak rate of 10.6% and Ryan et al identified 7,167 oesophagectomies with a trans-thoracic oesophagectomy leak rate of 9.8% and a trans-hiatal oesophagectomy leak rate of 12% (6, 7). However, until recently the definition of anastomotic leakage and gastric conduit necrosis have not been standardised across the surgical literature.

In 2015, the Esophagectomy Complications Consensus Group (ECCG) defined anastomotic leaks as full-thickness defects involving the oesophagus, anastomosis, staple line or conduit, irrespective of

the presentation or method of identification. In this classification, leaks were divided into three types based on management strategy. Type 1 leaks require no change in therapy, are treated medically or with dietary modification. Type 2 leaks require interventional but not surgical therapy (interventional radiology drain, stent, etc.) and type 3 leaks require surgical intervention.

Gastric conduit necrosis has also been classified by the ECCG Group, and this is when the gastric conduit becomes ischaemic and necrotic. Type 1 is focal gastric conduit necrosis which is identified endoscopically and is managed with increased monitoring and non-surgical therapy. Type 2 is focal gastric conduit necrosis which is identified endoscopically and not associated with a free anastomotic leak or conduit leak and is treated surgically, but did not require oesophageal diversion. Type 3 is extensive gastric conduit necrosis which is treated re-operation and resection of the necrotic stomach together with oesophageal diversion via cervical oesophagostomy.

The ECCG Group have recently published data on 2704 oesophageal resections operated on between January 2015 and December 2016. This data was from 24 high volume oesophageal units in 14 countries. The indication for resection was malignancy in 95.6%, neoadjuvant chemoradiation (46.1%) or neoadjuvant chemotherapy (29.5%) was given. In this dataset the anastomotic leak rate was 11.4% (95% CI 10.2-12.6) and the rate of conduit necrosis was (1.3% (95% CI 0.7-1.7) (16). In this study 52.1% of the oesophagectomies were performed open. In the ECCG study the 95% upper confidence limit for a major complications (Clavien Dindo grade 3 or worse) was 35.6%.

In a similar benchmarking study, from 13 high volume units over a 5 year period, outcomes from totally minimally invasive oesophagectomy (43.7% were 3 stage procedures) in low risk patients were defined. Anastomotic leakage in this cohort was 15.9%.

Numerous studies have advocated varying techniques comparing handsewn and mechanical options for anastomoses (17, 18). There is some evidence to show that a mechanical anastomosis using a linear stapler has a reduced leak rate and reduced stricture rate as compared to a handsewn anastomosis however results vary markedly between surgeons and units (19). There is evidence to

suggest that cervical anastomoses are associated with an increased leak rate as compared to thoracic anastomoses (7, 20).

Prompt recognition of anastomotic leakage can potentially speed clinical intervention and improve patient outcome. Early signs of anastomotic leakage include tachycardia, pyrexia, high white cell count, high CRP, delirium and cardiac arrhythmias, particularly atrial fibrillation. Late signs include bilious output from chest drain, acidosis, hypotension and septic shock. The clinical management of leaks are controversial and depend on the site of the leak, size of the defect, perfusion of the gastric conduit and the clinical status of the patient. Small contained anastomotic leaks can be managed conservatively without surgery, where patients are kept nil by mouth, and given antibiotics and nasogastric drainage. Leaks that are not localised or that cause greater systemic upset are generally considered to be those that require some form of active intervention such as radiological drainage or treatment with either endoluminal VAC therapy, covered oesophageal stenting or re-thoracotomy although there is little evidence of superiority of one technique over another. Large anastomotic leaks, especially if associated with severe sepsis or gastric conduit necrosis may require re-thoracotomy resection of the anastomosis and oesophageal diversion with cervical oesophagostomy. The main purpose of this international audit is to identify the incidence of leaks, see when they are diagnosed are diagnosed and how they are specifically managed.

An international multicentre audit will enable a large volume of patient data to be obtained over a short time period when changes in unit policies are likely to be minimised. It will potentially obtain a more general overview of the variations in practice across units and countries. Surgical access and anastomosis technique have been continued areas of disagreement amongst oesophago-gastric surgeons and their influence on mortality and morbidity has long been disputed. This audit seeks to provide up to date information in the international variations in practice.

Aim

Primary Audit Question

- 1- Quantify the incidence of anastomotic leak rate in an international multicentre audit which incorporates data from high and low volume centres and high and low income countries
- 2- Assess the variation in anastomotic leak rates internationally
- 3- Assess the relationship between anastomotic technique and optimal patient outcome – discharge home eating and drinking orally
- 4- Assess the relationship between anastomotic leak therapy and optimal patient outcome

Audit Standard

- 1- Anastomotic leak and conduit necrosis rate should be less than 13%
- 2- Major post-operative morbidity (Clavien Dindo Grade III or more) should be less than 35%.
- 3- 30 day mortality rate should be less than 5% and 90 day mortality rate should be less than 8%.

Primary Objective

The audit will aim to identify trends in patient factors and operative technique differences that may influence outcome. This in turn will allow for the formulation of more detailed research.

Key outcomes will include:

- Anastomotic Leak rate

OGAA Protocol 2018

- 30-day mortality

- 90-day mortality

- 30-day complication rate as set out in the International Consensus on Standardization of Data Collection for Complications Associated With Esophagectomy as defined by the Esophagectomy Complications Consensus Group (ECCG) (21)

- Length of stay

- Readmission

Methods

A global prospective audit of patients undergoing oesophagectomy over a 6 month period from April 2018 to October 2018. Patients will be followed up for 90 days after the date of surgical resection.

Registered units must include all patients undergoing oesophagectomy during the study period.

A 2 month pilot of 4 centres within the UK will be undertaken to finalise the detailed online case report forms. This will ensure that all relevant data is collected to achieve the goals of the audit.

Study Population

Inclusion Criteria

- All adult patients undergoing oesophagectomy for malignancy with an oesophagogastric anastomosis carried out during the study period.
- Any approach (2 stage Ivor Lewis, 3 stage McKeown, thoracoabdominal, trans-hiatal) using any combination of open, robotic or standard minimal access approaches)
- Malignant disease
- Elective (planned) resections.
- Thoracic and cervical anastomotic locations

Exclusion criteria

- Extended Total Gastrectomy
- Pharyngolaryngoesophagectomy
- Colonic interposition and small bowel jejunal interposition reconstructions
- Emergency resection
- Resections for benign disease

Patient identification

- Multidisciplinary team meetings
- Coordination with lead surgeon for oesophago-gastric cancer resections
- Coordination with Upper GI Cancer Specialist nursing services
- Review of theatre scheduling systems

Centre Eligibility

Any centre routinely performing elective oesophagectomies is eligible to join the audit. No restriction will be placed on global location or number of surgeons involved.

No restriction will be placed on the minimum number of oesophagectomies required to be enrolled in the audit.

Each unit will be required to register prior to the start date for data collection.

Each unit will be responsible for obtaining local hospital approval before commencement of the audit.

Each unit must ensure they have appropriate staff that will be able to ensure a >95% completeness of data entry before the closing date of the study.

Patient Follow Up

The study design aims to ensure that no additional patient follow up or intervention is required that would deviate from the normal patient journey.

For the purposes of accurate data entry follow up will require investigators to collate information from electronic and paper records. This will enable adequate analysis of the pre, intra and post-operative patient outcomes.

The data collection period will be for 90 days after the index operation involving the patient's first reconstruction.

Data Completion and Organisation

Data input will be via a dedicated encrypted server through the Research Electronic Data Capture (REDCap) web application. No patient identifiable information will be inputted into the database. REDCap will provide an ID number for each patient entered. Locally held records containing

corresponding REDCap ID numbers and local patient identifiers must be stored securely. This will facilitate patient data entry at different time points by different team members and enable cross checking of data entry by different team members to ensure accuracy of data collection.

An electronic REDCap “App” will be available for smart phones to enable data collection. Data will be held securely on the “App” and information can be uploaded to the central database when internet access is available. Printable data collection proformas will be made available to enable participants to record data as required that can be uploaded to REDCap when a computer/device is available.

Patient data will be entered into case report forms (CRFs) which are designed not to deviate from safe patient care. CRFs will only record patient events and not instigate any form of intervention.

Each unit will be able to register a maximum of 5 members who will be granted access to input unit data. Each unit will be required to have a lead auditor of Consultant grade (or equivalent, country dependent). Units may apply on an individual basis if they require additional team member registration.

Intra-operative detail must be entered by a surgeon present at the time of the operation. However if a nominated member of the audit is not present at the operation he/she must take instruction from a surgeon who was present at the time of the operation. This will minimise error and ensure accurate operative data recording that may be absent in operation note records. All other data such as demographics or outcomes may be inputted by any member of the audit team.

Missing data may be entered any time during the study period. Units with >5% missing data will be excluded from the study.

The Birmingham Surgical Trials Consortium, University of Birmingham, will host the REDCap system.

All data will be stored securely on encrypted and certified servers for a minimum of 5 years.

Data Collection Form

Please see appendix 1 for our detailed Data Collection / Case Report Form

Pre-operative variables, including patient demographics, age, gender, smoking and alcohol history, pre-operative blood results (Albumin, Haemoglobin, Creatinine) and co-morbidities will be collected here. These can be completed prior to the date of the operation if desired. We will also collect data on neoadjuvant therapy and pre-operative tumour stage.

Intra-operative variables, including the operation type, technique of the operation (open / laparoscopic / robotic), location of the anastomosis, type of anastomosis performed and any techniques to assess the anastomosis during the surgery will be documented here. Techniques to try to reduced anastomotic leakage, including wrapping the anastomosis in omentum or burying the anastomosis in the pleural will be collected.

Anaesthetic variables, we are also collecting information on single lung ventilation (double lumen tube / bronchial blocker), intra-operative infusion of fluids and blood and administration of vasopressors by bolus or infusion in the intra-operative period. We would also like to know the post procedure lactate level and whether the patient was extubated on the same day as the operation.

Post-operative / complications variables. These will mainly focus on the ECCG definitions of anastomotic leak and conduit necrosis (Appendix 3) and complications according to other organ systems (respiratory, cardiac, renal, chyle leak, etc). We will be assessing whether the anastomosis was formally assessed for integrity (by endoscopy / CT or contrast study) during the post-operative period. In addition, if an anastomotic leak or conduit necrosis does occur we will document the management strategy for the patient. This could change between the primary (first), secondary (second) and tertiary (third) management options – for example non-operative, operative (re-thoracotomy), radiological and endoscopic (stenting or endoVAC therapy) – in any order depending on what actually occurs to the patient. Final outcome data such as whether the patient was

discharged eating and drinking normally, total length of stay, 30 and 90 day mortality and readmission will also be collected.

Local Approvals

All data collected will measure current practice, with no changes made to normal treatment. As such, this study should be registered as an audit of current practice at each participating centre. It is the responsibility of the local team at each site to ensure that local audit approval (or equivalent) is completed for their centre. For example, surgeons and teams from other countries will have to abide by their local hospital / country approval process. Participating centres will be asked to confirm that they have gained formal approval at their site. Some international centres may require formal ethical approval to be obtained and some units may require individual patient consent. In the UK we have had confirmation that the project should be registered as an Audit (Appendix 2).

Authorship

A maximum of five investigators from each individual unit will be incorporated in this study as co-investigators. Investigators will be PubMed searchable and citable. The output from the study will be published under a single corporate authorship "Oesophagogastric Anastomosis Study Group, West Midlands Research Collaborative".

Pilot

A two month pilot held across four UK hospitals will be undertaken prior to the commencement of the full global audit. This will allow for potential adjustments to the investigation CRF proforma for a more comprehensive study.

Data Publication and Governance

Data will be published as pooled data. It is important to emphasise that no surgeon or unit specific data will be published. Local units may request their own specific data at the end of the study.

The “Oesophagogastric Anastomosis Study Group, West Midlands Research Collaborative” welcome the use of the data for further research. All requests will be assessed on an individual basis with a strong emphasis on safeguarding of data.

All subsequent publications using the dataset must recognise OAI and be published under the principals of shared authorship with a single corporate author.

International centres may require a data transfer agreement and this can be provided if required.

Funding

The Oesophagogastric Anastomosis Study Group currently has no specific funding, however the funding of the website (www.ogaa.org.uk) was kindly provided by funding from the Birmingham Oesophageal Cancer Patients Group which meets on a regular basis at the Queen Elizabeth Hospital, Birmingham.

Cohort size

We have estimated the number of eligible operations performed across Europe. Hospital Episode Statistics (HES) is a data warehouse containing details of all admissions at NHS hospitals in England. A HES database publication showed that over a ten year period between 2000 – 2010, an average of 1657 oesophagectomies were performed per year (22). The population of England is approximately 53 million. The population of Europe is approximately 739.2 million. Therefore if we accept the same rate $((1657/53,000,000) \times 739,200,000)$ there will be around 23,110 operations performed across Europe per year.

This prospective study will only pick-up a proportion of these patients, and this depends upon three factors: Penetration - the proportion of hospitals who sign up to recruit patients to the study across Europe; Pick-up - the proportion of the eligible patients at each centre are entered into the study; Study duration.

The following projection models have been estimated using various combinations of these three factors:

5% penetration; 80% pick-up 6 month recruitment = 924 cases

8% penetration; 90% pick-up 6 months recruitment = 1663 cases

10% penetration; 80% pick-up 6 month recruitment = 1848 cases

10% penetration; 90% pick-up 6 months recruitment = 2079 cases

20% penetration; 90% pick-up 6 month recruitment = 4159 cases

Caveats to these calculations include the variation in rates of oesophageal cancer and oesophagectomy in Europe and the intention that centres in other continents will also contribute to the study

Statistical analysis

The report of the audit will be prepared in accordance with the guidelines as set by the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) statement for observational studies and STROCCS (Strengthening the Reporting of Cohort Studies in Surgery). (1). Data will be collected and analysed in clinically relevant categories, and the Chi squared tests used to detect differences between groups. Missing data for predictor values will be replaced using the multiple imputation method to create five imputed datasets; all predictor and outcome variables will be entered into the predictive models for imputation.

Binary logistic regression modelling will be used. Multivariable models will be built to produce odds ratios (OR) to account for the impact of predictive variables when assessing outcomes (anastomotic leak). Variable selection will be based upon those which are statistically significant at univariable analysis, and those which are clinically significant but not statistically. Fixed, forced entry will be used to adjust the main outcome measure. The effect of interaction, and sequential removal of non-significant variables will be assessed using changes in Akaike information criterion for multilevel models, and p-values for multiply imputed fixed models. Finally, risk adjusted funnel plots will be produced to test the performance of individual (anonymised) centres for rates of anastomotic leak and other factors.

References

- 1- Agha, R. A., Borrelli, M. R., Vella-Baldacchino, M., Thavayogan, R., Orgill, D. P., Pagano, D., Pidgeon, T. E. (2017). The STROCCS statement: Strengthening the Reporting of Cohort Studies in Surgery. *International Journal of Surgery*, 46, 198–202.
- 2- Turrentine, Florence E., et al. "Morbidity, mortality, cost, and survival estimates of gastrointestinal anastomotic leaks." *Journal of the American College of Surgeons* 220.2 (2015): 195-206.
- 3- Low, Donald E. "Diagnosis and management of anastomotic leaks after esophagectomy." *Journal of Gastrointestinal Surgery* 15.8 (2011): 1319-1322.
- 4- Goense, L., van Dijk, W. A., Govaert, J. A., van Rossum, P. S. N., Ruurda, J. P., & van Hillegersberg, R. (2017). Hospital costs of complications after esophagectomy for cancer. *European Journal of Surgical Oncology*, 43(4), 696–702.
- 5- Markar, S., Gronnier, C., Duhamel, A., Mabrut, J.-Y., Bail, J.-P., Carrere, N., Mariette, C. (2015). The Impact of Severe Anastomotic Leak on Long-term Survival and Cancer Recurrence After Surgical Resection for Esophageal Malignancy. *Annals of Surgery*, 262(6), 972–980.
- 6- Ryan, Carrie E., et al. "Transthoracic anastomotic leak after esophagectomy: Current trends." *Annals of Surgical Oncology* 24.1 (2017): 281-290.
- 7- Kassis, Edmund S., et al. "Predictors of anastomotic leak after esophagectomy: an analysis of the society of thoracic surgeons general thoracic database." *The Annals of thoracic surgery* 96.6 (2013): 1919-1926.
- 8- Dent, B., et al. "Management and outcomes of anastomotic leaks after oesophagectomy." *British Journal of Surgery* 103.8 (2016): 1033-1038
- 9- Hu, Zhongwu, et al. "The diagnostic value of routine contrast esophagram in anastomotic leaks after esophagectomy." *World Journal of Surgery* (2017): 1-6.
- 10- Zehetner, Jörg, et al. "Intraoperative assessment of perfusion of the gastric graft and correlation with anastomotic leaks after esophagectomy." *Annals of surgery* 262.1 (2015): 74.
- 11- Bolton, John S., William C. Conway, and Abbas E. Abbas. "Planned delay of oral intake after esophagectomy reduces the cervical anastomotic leak rate and hospital length of stay." *Journal of Gastrointestinal Surgery* 18.2 (2014): 304-309.
- 12- Kanamori, J., et al. "Leak grading and percutaneous transanastomotic drainage for the treatment of cervical anastomotic leakage after esophagectomy." *Diseases of the Esophagus* 30.5 (2017): 1-7.
- 13- Roh, Simon, et al. "Role of Barium Swallow in Diagnosing Clinically Significant Anastomotic Leak following Esophagectomy." *The Korean journal of thoracic and cardiovascular surgery* 49.2 (2016): 99.
- 14- Guo, Juntang, et al. "Choice of therapeutic strategies in intrathoracic anastomotic leak following esophagectomy." *World journal of surgical oncology* 12.1 (2014): 402.
- 15- Perry, Yaron, et al. "Serial drain amylase can accurately detect anastomotic leak after esophagectomy and may facilitate early discharge." *The Annals of thoracic surgery* 100.6 (2015): 2041-2047.

- 16- Low, D. E., Kuppusamy, M. K., Alderson, D., Cecconello, I., Chang, A. C., Darling, G., Wijnhoven, B. P. L. (2017). Benchmarking Complications Associated with Esophagectomy. *Annals of Surgery*, 1.
- 17- Price, Theolyn N., et al. "A comprehensive review of anastomotic technique in 432 esophagectomies." *The Annals of thoracic surgery* 95.4 (2013): 1154-1161.
- 18- Markar, Sheraz R., et al. "Technical factors that affect anastomotic integrity following esophagectomy: systematic review and meta-analysis." *Annals of surgical oncology* 20.13 (2013): 4274-4281.
- 19- Deng, Xu-Feng, et al. "Hand-sewn vs linearly stapled esophagogastric anastomosis for esophageal cancer: a meta-analysis." *World Journal of Gastroenterology: WJG* 21.15 (2015): 4757.
- 20- Biere, S. S. A. Y., et al. "Cervical or thoracic anastomosis after esophagectomy for cancer: a systematic review and meta-analysis." *Digestive surgery* 28.1 (2011): 29-35.
- 21- Low, Donald E., et al. "International consensus on standardization of data collection for complications associated with esophagectomy: Esophagectomy Complications Consensus Group (ECCG)." *Annals of surgery* 262.2 (2015): 286-294.
- 22- Mamidanna, Ravikrishna, et al. "Surgeon volume and Cancer esophagectomy, gastrectomy, and pancreatectomy: a population-based study in England." *Annals of surgery* 263.4 (2016): 727-732.

Pre-Operative Data Collection

Gender	Male / Female
Age (in Years)	
ASA	1/2/3/4
Comorbidity (21-22) Ischaemic Heart Disease Cerebrovascular Disease Peripheral Vascular disease Diabetes Renal Disease Chronic Lung Disease Liver Disease	Yes / No Yes / No Yes / No Yes / No Yes / No Yes / No Yes / No
Smoking History	Never, Current, ex >6/52, ex <6/52
Height (cm) Weight (kg)	Automatic BMI Calculation
Pre-op bloods at start of surgery (or last recorded level, within previous 2 weeks) Albumin Haemoglobin eGFR (estimated Glomerular filtration rate)	____ g/L or mmol/L Absolute value in g/L to one decimal place [with pop-up converter to change from g/dL to mmol/L]
Malignancy details Tumour type Location of tumour Neo-adjuvant therapy Overall Pre-operative staging If Radiotherapy give pre-op	Adeno / SCC / Other Upper / Mid / Siewert 1 / 2 / 3 None / Chemotherapy / Chemoradiotherapy TNM 7 th Total Gy _____ Did the radiotherapy field include the gastric fundus – yes / no
Pre-operative nutritional support	None Oral Supplements Enteral Nutrition via NJ/NG/PEG/Jej etc TPN
Pre-operative gastric ischaemic preconditioning performed *	Yes / No

* This is when laparoscopy and division of the left gastric vessels +/- short gastric vessels are performed prior to oesophagectomy under a separate anaesthetic

Intra Operative Data Collection

Training operation	Yes / No
Trainee performed abdominal phase	Yes / No
Trainee performed chest dissection	Yes / No
Trainee performed anastomosis	Yes / No
Abdominal phase	Lap / Open / Lap Converted to open / Robotic
Thoracic phase	Thorascopic / Open Right Chest / Open Left chest or thoracoabdominal / Thorascopic converted to open / Trans-hiatal / Robotic
Lymphadenectomy	Abdominal only Abdominal and Thoracic (2 field) Abdominal / Thoracic / Neck (3 field)
Gastric Tube	Whole Stomach, Wide Gastric Tube > 5cm, Thin Gastric Tube < 5cm
Anastomosis level	Neck / Chest above Azygous / At Azygous / Below Azygous / Anastomosis not performed
Anastomotic configuration	End to End Side to End Side to Side
Anastomosis technique	
Handsewn	Single layer / Two layer Interrupted / Continuous
Circular stapler	Type- CEA / CDH / other- please specify Head diameter (mm)- please specify OrVil (25mm)
Orringer style anastomosis (linear stapled and sutured)	Yes / No
Was the anastomosis wrapped or covered in omentum	Yes / No
Was the anastomosis buried in mediastinal pleura	Yes / No
Was the anastomosis tested for integrity	Not performed / NG Air Leak Test / Intra-op Endoscopy / Methylene Blue / Indigocyanine green (IGC) assessment / Other method
Nutritional Feeding Access	None / Feeding Jejunostomy / Nasojejunal tube
Procedures on the Pylorus	None/ Pyloromyotomy / Pyloroplasty / Botox / Dilatation / other
Intra-op complications	Yes / No

	Please specify
Total Operative duration (mins)	

Anaesthetic Data Collection

Single Lung Ventilation	Yes / No If Yes – Double Lumen Tube or Bronchial Blocker If Yes - Duration of One Lung Ventilation (mins)
Intra-operative vasopressor support required (For example Noradrenaline, Metaraminol, Ephedrine or phenylephedrine etc)	Yes – bolus Yes – continuous infusion No
Total IV Fluid (mls) given intra-operatively	_____ mls crystalloid _____ mls colloid
Intra-operative blood transfusion	Yes / No If Yes - Number of units transfused _____
Analgesia technique	Epidural Thoracic paravertebral block Intra-thecal Morphine Patient Controlled Analgesia (PCA) Ketamine Abdominal pain catheter
Lactate Level immediately postoperative	_____ mmol/L
Was the patient extubated the same day as resectional surgery?	Yes / No

Post Operative Data Collection

Re-intubated	Yes / No
Return to ICU	Yes / No
Return to theatre	Yes / No
Was assessment of anastomosis performed in the post op period? Endoscopy Plain Film Contrast Swallow CT Contrast Swallow Other	Yes / No Yes / No Yes / No Please specify
What day post operatively did this occur	Post-op Day _____
Post Operative Complications Anastomotic leak No. of days after surgery leak was diagnosed Conduit Necrosis No. of days after surgery conduit necrosis was diagnosed Chyle Leak Pneumonia Diaphragmatic hernia Feeding jejunostomy complication Cardiac complication DVT or PE Other significant complication	Yes / No / Grade 1 / 2 / 3 No days _____ Yes / No / Grade 1 / 2 / 3 No days _____ Yes / No / Grade 1 / 2 / 3 Yes / No Yes / No Yes / No Yes / No Yes / No Yes / No Details _____
Primary Treatment of leak/conduit necrosis Post-operative day of start of treatment _____ Primary treatment successful – Yes / No	Non-operative management – Yes / No Radiological drainage – Yes / No Oesophageal stenting – Yes / No If Stent – Covered Plastic / Covered Metal Successful / Unsuccessful

	<p>Complications from Stent – Displacement / Erosion / Failure to Occlude Leak / Other Re-stented Total no of stents used</p> <p>Endoluminal VAC therapy – Yes / No Total no of EndoSponge changes Re-operation:</p> <p>Opening of Neck Wound</p> <p>Minimal Access or Open Thoracotomy</p> <p>Washout only / Anastomotic Repair / Reformation of the Anastomosis / T-Tube / Intercostal or muscle flap repair / Disconnection and cervical oesophagostomy</p>
<p>Secondary Leak Treatment of leak/conduit necrosis</p> <p>Post-operative day of start of treatment _____</p> <p>Secondary treatment successful – Yes / No</p>	<p>Non-operative management – Yes / No</p> <p>Radiological drainage – Yes / No</p> <p>Oesophageal stenting – Yes / No</p> <p>If Stent – Covered Plastic / Covered Metal Successful / Unsuccessful Complications from Stent – Displacement / Erosion / Failure to Occlude Leak / Other Re-stented Total no of stents used</p> <p>Endoluminal VAC therapy – Yes / No Total no of EndoSponge changes Re-operation:</p> <p>Opening of Neck Wound</p> <p>Minimal Access or Open Thoracotomy</p> <p>Washout only / Anastomotic Repair / Reformation of the Anastomosis / T-Tube / Intercostal or muscle flap repair / Disconnection and cervical oesophagostomy</p>
<p>Total Length of stay of hospital stay (Days)</p>	
<p>Total length of ICU and HDU stay (Days)</p>	
<p>Was the patient discharged from hospital eating and drinking?</p>	<p>Yes / No</p>
<p>Final Histology (23-24)</p>	

<p>T stage No Nodes examined No Nodes positive for malignancy</p> <p>Surgical Margins</p> <p>M stage</p>	<p>Complete path response / HGD / 1 / 2 / 3 / 4 No nodes _____ No nodes _____</p> <p>Proximal margin – clear / involved (<1mm) Distal margin – clear / involved (<1mm) Circumferential / radial margin – clear / involved (<1mm)</p> <p>0/1</p>
<p>In hospital post-operative death Within 30 days of surgery? Within 90 days of surgery?</p>	<p>Yes / No Yes / No Yes / No</p>
<p>Out of hospital post-operative death Within 30 days of surgery? Within 90 days of surgery?</p>	<p>Yes / No Yes / No Yes / No</p>
<p>30 day readmission</p>	<p>Yes / No</p>

Unit Questionnaire

Number of consultant surgeons performing oesophagectomy	Total No.
Number of oesophagectomies performed between Jan 2015 and Dec 2016	
Speciality of Surgeons	Thoracic / Oesophagogastric / General Surgeon / Surgical Oncologist
Size of institution	Total number of beds Total number of ICU beds
24 hour on call rota for oesophageal emergencies	24hour / 9-5 / none
24 hour on call availability for interventional radiology	24hour / 9-5 / none
24 hour access to emergency theatre	24hour / 9-5 / none
Where do oesophagectomy patients routinely go post-operatively	Ward HDU ICU Dedicated GI HDU
ERAS protocol for oesophagectomy patients	Yes / No
ERAS nurse Physio input	Yes / No Nil dedicated / Daily / Twice daily
Does your unit perform gastric ischaemic preconditioning?	Yes – Routinely Yes – Selectively No If Yes – how many days prior to surgery
Does your unit have an agreed approach to oesophagectomy for lower 1/3 adenocarcinoma?	No Yes Open Right Transthoracic Oesophagectomy Open Left thoracoabdominal oesophagectomy Open Transhiatal Oesophagectomy Hybrid Transthoracic Oesophagectomy (Lap abdominal mobilisation) 2 stage Minimal Access Oesophagectomy 3 stage Minimal Access Oesophagectomy Robotic Oesophagectomy Other
Does your unit have an agreed technique to perform intra-thoracic anastomosis?	No Yes Handsew Circular Stapled OrVil

	<p>Stapled side to side with suturing (Orringer style) Other</p>
Does your unit have access to Indigo-Cyanine Green assessment of the anastomosis or gastric conduit?	Yes / No
Does your unit have a policy of performing routine post-operative assessment of the anastomosis?	<p>No</p> <p>Yes – Barium or Water Soluble Contrast Swallow</p> <p>Yes – Endoscopy</p> <p>Yes – CT</p> <p>If your unit routinely assess the anastomosis in the post-operative period, what day is this generally performed?</p> <p>Postop Day _____</p>
Does your unit have access to following for the treatment of oesophageal anastomotic leak?	<p>TPN – Yes / No</p> <p>Endoscopic Clips – Yes / No</p> <p>Endoscopic or radiologically placed covered oesophageal stents – Yes / No</p> <p>EndoVAC / Endosponge therapy – Yes / No</p> <p>Interventional guided drainage of abdominal or thoracic collections – Yes / No</p>

Appendix 1 - How to register this audit

Every hospital has an audit department which should be able to advise on the information required to register the project. Please contact them well in advance to ensure all the paper work is correct (we would recommend at least one month prior to the study commencing).

At Trust level:

1. Identify a PI (Primary Investigator) at each trust – this is a Consultant who agrees to support the study.
2. Create a team of Consultants/ surgical registrars.
3. Contact your hospital's Clinical Audit Department preferably by email
 - a. They will provide you with a standard audit form to complete, via email or from the intranet
 - b. You can copy and paste from this protocol
 - c. Ensure that the audit department know that this is part of a larger project and that you will send anonymised data for central collation via secure nhs.net email addresses. This will involve gaining permission from the Trust's Caldicott Guardian if based in the UK.
4. Once the form is completed, you may need to ask your supervising consultant to sign it.
5. Form submission
 - a. Submit the form and protocol to the Audit Department as soon as possible.
6. Email form to OGanastomosisaudit@gmail.com to register your interest.

Appendix 2 - Health Research Authority Tool UK



Is my study research?

I To print your result with title and IRAS Project ID please enter your details below:

Title of your research:

Oesophageal Anastomosis Investigation

IRAS Project ID (if available):

You selected:

- 'No' - Are the participants in your study randomised to different groups?
- 'No' - Does your study protocol demand changing treatment/ patient care from accepted standards for any of the patients involved?
- 'No' - Are your findings going to be generalisable?

Your study would NOT be considered Research by the NHS.

You may still need other approvals.

Researchers requiring further advice (e.g. those not confident with the outcome of this tool) should contact their R&D office or sponsor in the first instance, or the [HRA](#) to discuss your study. If contacting the HRA for advice, do this by sending an outline of the project (maximum one page), summarising its purpose, methodology, type of participant and planned location as well as a copy of this results page and a summary of the aspects of the decision(s) that you need further advice on to the HRA Queries Line at HRA.Queries@nhs.net.

Appendix 3 - Grading Oesophageal Complications

Anastomotic Leak

Defined as: Full thickness GI defect involving oesophagus, anastomosis, staple line, or conduit irrespective of presentation or method of identification

Type I: Local defect requiring no change in therapy or treated medically or with dietary modification

Type II: Localized defect requiring interventional but not surgical therapy, for example, interventional radiology drain, stent or bedside opening, and packing of incision

Type III: Localized defect requiring surgical therapy

Conduit Necrosis

Type I: Conduit necrosis focal Identified endoscopically

Treatment — Additional monitoring or non-surgical therapy

Type II: Conduit necrosis focal Identified endoscopically and not associated with free anastomotic or conduit leak

Treatment — Surgical therapy not involving esophageal diversion

Type III: Conduit necrosis extensive

Treatment — Treated with conduit resection with diversion

Low, Donald E., et al. "International consensus on standardization of data collection for complications associated with esophagectomy: Esophagectomy Complications Consensus Group (ECCG)." *Annals of surgery* 262.2 (2015): 286-294

Appendix 4 – TNM Staging (7th Edition, 23-24)

Primary Tumour (T)

TX	Primary tumour cannot be assessed
T0	No evidence of primary tumour
Tis	High-grade dysplasia
T1	Tumour invades lamina propria, muscularis mucosae, or submucosa
T1a	Tumour invades lamina propria or muscularis mucosae
T1b	Tumour invades submucosa
T2	Tumour invades muscularis propria
T3	Tumour invades adventitia
T4	Tumour invades adjacent structures
T4a	Resectable tumour invading pleura, pericardium, or diaphragm
T4b	Unresectable tumour invading other adjacent structures, such as the aorta, vertebral body, and trachea

Regional lymph nodes (N)

NX	Regional lymph node(s) cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in 1-2 regional lymph nodes
N2	Metastasis in 3-6 regional lymph nodes
N3	Metastasis in 7 or more regional lymph nodes

Distant metastasis (M)

M0	No distant metastasis
M1	Distant metastasis